arXiv:1208.2857v2 [quant-ph] 24 Sep 2012

On Requirement of Dissonance in Assisted Optimal State Discrimination

Fu-Lin Zhang,[1,*] Jing-Ling Chen,[2,3,†] L. C. Kwek,[3,4] and Vlatko Vedral[3,5,6]
[1] *Physics Department, School of Science, Tianjin University, Tianjin 300072, China*
[2] *Theoretical Physics Division, Chern Institute of Mathematics,*
Nankai University, Tianjin, 300071, P. R.China
[3] *Centre for Quantum Technologies, National University of Singapore, 3 Science Drive 2, Singapore 117543*
[4] *National Institute of Education and Institute of Advanced Studies,*
Nanyang Technological University, 1 Nanyang Walk, Singapore 637616
[5] *Clarendon Laboratory, University of Oxford, Parks Road, Oxford OX1 3PU, United Kingdom*
[6] *Department of Physics, National University of Singapore, 2 Science Drive 3, Singapore 117542*
(Dated: September 25, 2012)

We reexamined the roles of entanglement, discord, and dissonance needed in the procedure of optimal quantum state discrimination assisted by an auxiliary system. We found that entanglement between the principal qubit and the ancilla is completely unnecessary, which can be zero in arbitrary case by adjusting the parameters in the more general joint unitary transformation without affecting the success probability. This result strengthens the argument that quantum dissonance indeed plays as a key ingredient in the assisted optimal state discrimination other than entanglement.

PACS numbers: 03.65.Ta, 03.67.Mn, 42.50.Dv

Introduction.—As an essential distinction between quantum mechanics and classical mechanics, quantum coherent superposition leads to nonclassical correlations in the composite quantum systems: quantum entanglement [1], Bell nonlocality [2] and quantum discord [3, 4]. Quantum entanglement has been widely studied from various perspectives, and it has been found to be a useful resource to demonstrate the superiority of quantum information processing. For instances, entangled states are regarded as key resources in some quantum information tasks, such as teleportation and superdense coding [5]; and Bell nonlocality, characterized by violation of Bell inequalities, may guarantee the security of quantum cryptography [6].

In contrast to entanglement, the so-called quantum discord is presented as the quantum part (opposite to the classical one) of the total correlation in a bipartite system. Recent developments of deterministic quantum computation with one qubit (DQC1) [7, 8] indicate that a quantum algorithm for computing the trace of a unitary matrix can surpass the corresponding classical algorithm in terms of speedup even in the absence of quantum entanglement. This surprising result has generated wide interest in recent years, leading to many good works on quantum discord and its relations with other correlations. For instance, it has been shown that it is possible to have an operational interpretation in terms of the quantum state merging protocol [9, 10], and that it serves as the entanglement generated in an activation protocol [11] or in a measurement process [12]. In a unified view of quantum correlations based on the relative entropy [13], a new measure, quantum dissonance, is regarded as the nonclassical correlations in which entanglement has been excluded. For a separable state (with zero entanglement), its quantum dissonance is just equal to its quantum discord.

It is always a significant subject to investigate the roles of nonclassical correlations in quantum information processing. Recently the quantum algorithm of DQC1 has been widely accepted as the first example for which quantum dissonance plays as a key ingredient other than entanglement. This gives rise to a natural question: Can quantum dissonance also serve as a similar key resource in some other quantum information tasks? The answer is affirmative. Very recently, in an illuminating work of Roa, Retamal and Alid-Vaccarezza (RRA), they studied the roles of entanglement, discord, and dissonance needed for performing unambiguous quantum state discrimination assisted by an auxiliary qubit [14], which is a fundamental problem in both quantum mechanics and quantum information theory [15–19]. The RRA protocol in general requires both entanglement and discord in the *assisted optimal state discrimination* (AOSD), except one particular case with equal *a priori* probabilities (i.e., $p_+ = p_-$, where p_+ and p_- are probabilities prepared for the two states $|\psi_+\rangle$ and $|\psi_-\rangle$, respectively). In this specific cases, the entanglement of the state of system-ancilla qubits is absent but its discord is nonzero, and hence the unambiguous state discrimination protocol is implemented successfully only by quantum dissonance.

Clearly, the RRA protocol can be viewed as the second example for which dissonance plays as a key ingredient for AOSD other than entanglement. However, the original RRA protocol belongs to a two-parameter AOSD protocol, thus this prevents them from observing that in general entanglement is unnecessary for the AOSD. In this Letter, we reexamine the roles of correlations in the AOSD and extend the subject further. We find that in one case, only dissonance is required for the AOSD, and in another case, dissonance can be zero and the classical state may accomplish the AOSD. In both cases, entanglement is completely not required.

The Generalized RRA Protocol.—Let us consider the same model as in [14], in which a qubit is randomly prepared in one of the two nonorthogonal states $|\psi_+\rangle$ or $|\psi_-\rangle$ with *a priori* probabilities p_+ and p_-, respectively, here $p_+ + p_- = 1$. To discriminate the two states $|\psi_+\rangle$ or $|\psi_-\rangle$ unambiguously, the system is coupled to an auxiliary qubit A, which is prepared in a known initial pure state $|k\rangle_a$. Under a joint unitary transformation $\mathcal{U}$ between the system and the ancilla, one obtains [20]

$$\mathcal{U}\,|\psi_+\rangle|k\rangle_a = \sqrt{1-|\alpha_+|^2}\;|0\rangle|0\rangle_a + \alpha_+|\Phi\rangle|1\rangle_a, \qquad (1a)$$

$$\mathcal{U}\,|\psi_-\rangle|k\rangle_a = \sqrt{1-|\alpha_-|^2}\;|1\rangle|0\rangle_a + \alpha_-|\Phi\rangle|1\rangle_a, \qquad (1b)$$

where $|\Phi\rangle = \cos\beta|0\rangle + \sin\beta e^{i\delta}|1\rangle$, $\{|0\rangle,|1\rangle\}$ and $\{|0\rangle_a,|1\rangle_a\}$ are the orthonormal bases for the system and the ancilla, respectively. The probability amplitudes α_+ and α_- satisfy $\alpha_+^*\alpha_- = \alpha$, where $\alpha = \langle\psi_+|\psi_-\rangle = |\alpha|e^{i\theta}$ is the *priori* fixed overlap between the two nonorthogonal states. Note that Eq. (1) contains four independent parameters $\{\alpha,\alpha_+,\beta,\delta\}$, thus the generalized RRA protocol considered here is a four-parameter protocol.

The state for the system-ancilla qubits is given by

$$\begin{aligned}\rho_{SA} &= p_+\,\mathcal{U}\left(|\psi_+\rangle\langle\psi_+|\otimes|k\rangle_a\langle k|\right)\mathcal{U}^\dagger \\ &\quad + p_-\,\mathcal{U}\left(|\psi_-\rangle\langle\psi_-|\otimes|k\rangle_a\langle k|\right)\mathcal{U}^\dagger, \end{aligned}\qquad (2)$$

which depends on β and δ, and generally is not equivalent to the corresponding one in [14] under local unitary transformations. Similar to [14], the recognition is successful when the ancilla collapses to $|0\rangle_a$. Therefore, the probability of success is achieved as

$$\begin{aligned}P_{\rm suc} &= {\rm Tr}[(\mathbb{1}_s\otimes|0\rangle_a\langle 0|)\rho_{SA}] \\ &= p_+(1-|\alpha_+|^2) + p_-(1-|\alpha_-|^2), \end{aligned}\qquad (3)$$

where $\mathbb{1}_s$ is the unit matrix for the system qubit. Without loss of generality as well as for simplicity, let us assume that $p_+ \le p_-$ and denote $\bar{\alpha} = \sqrt{p_+/p_-}$. The analysis of the optimal success probability can be divided into two different cases: (i) $|\alpha| < \bar{\alpha}$, $P_{\rm suc}$ is attained for $|\alpha_+| = \sqrt[4]{p_-/p_+}\sqrt{|\alpha|}$; (ii) $\bar{\alpha} \le |\alpha| \le 1$, $P_{\rm suc}$ is attained for $|\alpha_+| = 1$ (or equivalently $|\alpha_-| = |\alpha|$). Explicitly one has

$$P_{\rm suc,max} = \quad 1 - 2\sqrt{p_+p_-}|\alpha|, \quad \text{for case(i)}, \qquad (4a)$$

$$P_{\rm suc,max} = \quad (1-|\alpha|^2)p_-, \quad \text{for case(ii)}. \qquad (4b)$$

Note that $P_{\rm suc}$ is independent of the parameters β and δ, thus Eqs. (3) and (4) recover the results of [14].

Now we are ready to reexamine the roles of correlations playing in the *assisted optimal state discrimination.* To do this, let us calculate the entanglement degree of ρ_{SA}: the analytical expression for its concurrence [21] is given by

$$\begin{aligned}\mathcal{C}(\rho_{SA}) &= 2\Big[\mathcal{Y}_+^2\sin^2\beta + \mathcal{Y}_-^2\cos^2\beta \\ &\quad -2\mathcal{Y}_+\mathcal{Y}_-\sin\beta\cos\beta\cos(\theta+\delta)\Big]^{1/2}, \end{aligned}\qquad (5)$$

with $\mathcal{Y}_\pm = \sqrt{1-|\alpha_\pm|^2}\;|\alpha_\pm|p_\pm$. When $\beta = \pi/4$ and $\delta = 0$, Eq. (5) returns to the result in [14].

Let us impose the constraint $\mathcal{C}(\rho_{SA}) = 0$ for any α, α_+ and p_+. It is then easy to have the following solution

$$\delta = -\theta, \quad \beta = \arctan(\mathcal{Y}_-/\mathcal{Y}_+). \qquad (6)$$

Based on Eq. (6), state (2) is a separable state as

$$\rho_{SA} = |\eta_1\rangle\langle\eta_1|\otimes|0\rangle_a\langle 0| + |\Phi\rangle\langle\Phi|\otimes|\eta_2\rangle_a\langle\eta_2|, \qquad (7)$$

where $|\eta_1\rangle$ and $|\eta_2\rangle_a$ are two unnormalized states as

$$\begin{aligned}|\eta_1\rangle &= \frac{\sqrt{p_+p_-}}{\mathcal{Z}}\left(\sqrt{1-|\alpha_+|^2}\alpha_-|0\rangle - \sqrt{1-|\alpha_-|^2}\alpha_+|1\rangle\right), \\ |\eta_2\rangle_a &= \frac{\sqrt{\mathcal{Y}_+^2+\mathcal{Y}_-^2}}{\mathcal{Z}}\;|0\rangle_a + \frac{\mathcal{Z}\alpha_+}{|\alpha_+|}\;|1\rangle_a. \end{aligned}\qquad (8)$$

where $\mathcal{Z} = \sqrt{p_+|\alpha_+|^2 + p_-|\alpha_-|^2}$.

Note that the rank of the state (2) is two, thus its discord can be derived analytically by using the Koashi-Winter identity [22]. Namely, state (2) can be considered as a reduced state of the following tripartite pure state

$$|\Psi\rangle = \sqrt{p_+}\;(\mathcal{U}|\psi_+\rangle|k\rangle_a)|0\rangle_c + \sqrt{p_-}\;(\mathcal{U}|\psi_-\rangle|k\rangle_a)|1\rangle_c, \qquad (9)$$

where a fictitious qubit C is introduced. Then the discord can be written as $D(\rho_{SA}) = S(\rho_A) - S(\rho_{SA}) + E(\rho_{SC}) = S(\rho_A) - S(\rho_C) + E(\rho_{SC})$, where $S(\rho)$ is the von Neumann entropy, $E(\rho_{SC})$ is the entanglement of formation [21] between the principal system and the fictitious qubit C. Explicitly, the discord can be calculated as

$$D(\rho_{SA}) = \mathcal{H}(\tau_A) - \mathcal{H}(\tau_C) + \mathcal{H}(\tau_{SC}), \qquad (10)$$

where $\mathcal{H}(x) = -\frac{1+\sqrt{1-x}}{2}\ln\frac{1+\sqrt{1-x}}{2} - \frac{1-\sqrt{1-x}}{2}\ln\frac{1-\sqrt{1-x}}{2}$, τ_A is the tangle between A and SC, τ_C is the tangle between C and SA, and $\tau_{SC} = \mathcal{C}^2(\rho_{SC})$ is the concurrence between S and C in the state ρ_{SC}. One can obtains

$$\tau_A = \tau_{SA} + 4p_+p_-(|\alpha_+|^2 + |\alpha_-|^2 - 2|\alpha|^2), \qquad (11a)$$

$$\tau_C = 4p_+p_-(1-|\alpha|^2), \qquad (11b)$$

$$\tau_{SC} = \tau_S - \tau_{SA} - \tau(|\Psi\rangle), \qquad (11c)$$

with τ_S the tangle between S and AC, $\tau_{SA} = \mathcal{C}^2(\rho_{SA})$,



FIG. 1: Quantum dissonance in the AOSD. We plot the dissonance versus $|\alpha|$, for $p_+ = 1/2$ (solid line), $1/4$ (dashed line), and $1/8$ (dotted line). Dissonance is greater than zero for case (i), and is zero for case (ii). The critical point for $D(\rho_{SA}) = 0$ occurs at $|\alpha| = \bar{\alpha}$.

and $\tau(|\Psi\rangle)$ the three-tangle [23]. Explicitly we have

$$\begin{aligned}\tau_S &= 4\{(p_-|\alpha_-|^2 + p_+|\alpha_+|^2)[p_-(1-|\alpha_-|^2)\cos^2\beta \\ &\quad + p_+(1-|\alpha_+|^2)\sin^2\beta] \\ &\quad + p_+p_-(1-|\alpha_+|^2)(1-|\alpha_-|^2)\}, \end{aligned} \qquad (12)$$

$$\begin{aligned}\tau(|\Psi\rangle) &= 4p_+p_-|\sqrt{1-|\alpha_-|^2}\alpha_+\cos\beta \\ &\quad + \sqrt{1-|\alpha_+|^2}\alpha_-\sin\beta e^{i\delta}|^2. \end{aligned} \qquad (13)$$

It can be checked that $\tau(|\Psi\rangle)$ is independent on θ, if we substitute $\delta = -\theta$ and $\alpha_- = \alpha\alpha_+/|\alpha_+|^2$ into Eq. (13).

Dissonance for cases (i) and (ii).— For case (i), after substituting $|\alpha_+| = \sqrt[4]{p_-/p_+}\sqrt{|\alpha|}$, $p_- = 1 - p_+$, and Eqs. (6)(11)(12)(13) into Eq. (10), one then has the analytical expression for the dissonance, which depends only on $|\alpha|$ and p_+. In Fig. 1, we plot the curves of the dissonance versus $|\alpha|$ for $p_+ = 1/2, 1/4, 1/8$, respectively (see the curves with $D(\rho_{SA}) > 0$). For case (ii), because $|\alpha_+| = 1$, one has $\beta = \pi/2$ and thus the state ρ_{SA} becomes

$$\rho_{SA} = |1\rangle\langle 1| \otimes \rho_a, \qquad (14)$$

with $\rho_a = p_+|1\rangle\langle 1| + p_-|\mu\rangle_a\langle\mu|$, $|\mu\rangle_a = \sqrt{1-|\alpha|^2}\,|0\rangle_a + \alpha_- e^{i\delta}|1\rangle_a$. Obviously the state (14) is a direct-product state hence its dissonance is zero. In Fig. 1, for case (ii), we also plot the curves of dissonance versus $|\alpha|$ for the same p_+'s (see the curves with $D(\rho_{SA}) = 0$). Fig. 1 indicates that dissonance plays a key ingredient for AOSD other than entanglement for case (i), and the classical state may accomplish the task of AOSD for case (ii).

Geometric Picture.— It can be observed that the analysis of the optimal success probability $P_{\mathrm{suc,max}}$ in Eq. (4) was divided into two regions: $|\alpha| < \bar{\alpha}$ and $|\alpha| \geq \bar{\alpha}$. Here based on the positive-operator-valued measure (POVM) strategy [18, 19], we would like to provide a geometric picture to show why $P_{\mathrm{suc,max}}$ looks like that. Since the success probability, the concurrence and the discord of state ρ_{SA} are all independent on the phase θ of α (hence one can simply set $\theta = 0$), one can regard the states $|\psi_\pm\rangle$ as two unit vectors in $\mathbb{R}^2$ with the angle $\gamma = \arccos|\alpha|$ between them. The square roots of the *priori* probabilities, i.e., $\sqrt{p_+}$ and $\sqrt{p_-}$, play the roles of quantum wave amplitudes, which can be seen from the construction in Eq. (9). In Fig. 2, we plot two vectors $\overrightarrow{OA}$ and $\overrightarrow{OB}$ with $\angle BOA = \gamma$ to denote $\sqrt{p_+}|\psi_+\rangle$ and $\sqrt{p_-}|\psi_-\rangle$, respectively. The two POVM elements that identify the states $\sqrt{p_\pm}|\psi_\pm\rangle$ can be constructed as $\Pi_\pm = r_\pm|\tilde{\psi}_\pm\rangle\langle\tilde{\psi}_\pm|/(\langle\tilde{\psi}_\pm|\tilde{\psi}_\pm\rangle)$, with $r_\pm \geq 0$, and $|\tilde{\psi}_\pm\rangle = |\psi_\pm\rangle - |\psi_\mp\rangle\langle\psi_\mp|\psi_\pm\rangle$ the components of $|\psi_\pm\rangle$ orthogonal to $|\psi_\mp\rangle$. The vectors $\overrightarrow{AC}$ and $\overrightarrow{BD}$ correspond to the unnormalized states $|\tilde{\psi}_\pm\rangle$ with the coefficients $-\sqrt{p_\pm}$. The third POVM element giving the inconclusive result is $\Pi_0 = \mathbb{1}_s - \Pi_+ - \Pi_-$. The elements $\Pi_{\pm,0}$ are required to be positive, this is the constraint on the POVM strategy. Then the probability of successful discrimination is $P_{POVM} = (r_+p_+ + r_-p_-)(1-|\alpha|^2)$, and it can be expressed as


B
$-r_+\sqrt{p_+}\,|\tilde{\psi}_+\rangle$
E
$\sqrt{p_-}\,|\psi_-\rangle$
C
$-\sqrt{p_-}\,|\tilde{\psi}_-\rangle$
$-r_-\sqrt{p_-}\,|\tilde{\psi}_-\rangle$
$-\sqrt{p_+}\,|\tilde{\psi}_+\rangle$
γ
O
D
$\sqrt{p_+}\,|\psi_+\rangle$
A

FIG. 2: Geometric picture for optimal success probability based on POVM strategy. The sides $|OA| = \sqrt{p_+}$, $|OB| = \sqrt{p_-}$, the angle $\gamma = \arccos|\alpha|$, and $AC \perp OB$, $BD \perp OA$, $EB \perp OB$, $EA \perp OA$. For $|\alpha| < \bar{\alpha}$, the point E locates inside of the angle $\angle AOB$; for $|\alpha| \geq \bar{\alpha}$, the point E coincides with the point B for $p_+ < p_-$ (or A for $p_+ > p_-$).

$$P_{POVM} = (\overrightarrow{OA} - \overrightarrow{OB}) \cdot (r_-\overrightarrow{BD} - r_+\overrightarrow{AC}). \qquad (15)$$

When $|\alpha| < \bar{\alpha}$, the optimal P_{POVM} is attained at $r_\pm = (1 - \cos\gamma\sqrt{p_\mp/p_\pm})/\sin^2\gamma$. Then, the vectors $r_-\overrightarrow{BD} = \overrightarrow{EA}$ and $r_+\overrightarrow{AC} = \overrightarrow{EB}$ precisely, where E is the intersection point of AE and BE (see Fig. 2). The maximum value of P_{POVM} is the square of $|AB|$, nanmely $P_{POVM} = |AB|^2 = 1 - 2\sqrt{p_+p_-}|\alpha|$, which recovers Eq. (4a). When $|\alpha| = \bar{\alpha}$, the point E coincides with B



FIG. 3: Necessary criterion for requiring dissonance in AOSD based on linear entropy or purity. The linear entropy reads $\mathcal{S}_L(\rho) = 2 - 2\,\mathrm{Tr}\rho^2 = 4p_+p_-(1-|\alpha|^2)$, and the purity $\mathcal{P}(\rho) = 1 - \mathcal{S}_L(\rho)$. For a given amount of $|\alpha|$, when $\mathcal{S}_L(\rho) \leq 4(1-|\alpha|^2)|\alpha|^2/(1+|\alpha|^2)^2$, the dissonance for the AOSD is zero, see the region below the dashed line. $\mathcal{S}_L^{\max}(\rho) = 1/2$ when $|\alpha| = 1/\sqrt{3}$. Namely, if $\mathcal{S}_L(\rho) > 1/2$, then the dissonance is necessarily required for the AOSD.

for $p_+ < p_-$ (or A for $p_+ > p_-$), for the optimal P_{POVM} one has $r_- = 1$ (or $r_+ = 1$) and $P_{POVM} = p_-(1-|\alpha|^2)$. When $|\alpha| > \bar{\alpha}$, for $p_+ < p_-$, E is outside of the angle $\angle AOB$ and $\overrightarrow{EB}$ is opposite to $\overrightarrow{AC}$, which does not provide a physically realizable value of r_+. The optimal P_{POVM} strategy occurs at $r_- = 1$ and $r_+ = 0$ (i.e., point E coincides with point B), one then has $P_{POVM} = -\overrightarrow{OB} \cdot \overrightarrow{BD} = p_-(1-|\alpha|^2)$, which recovers Eq. (4b).

Conclusion and Discussion.—In summary, we have studied the generalized RRA protocol for optimal state discrimination assisted by an auxiliary qubit and found a significantly less stringent requirement on quantum correlations than the original result in [14]. The entanglement between the principal qubit and the ancilla is completely unnecessary, and it can be zero in the arbitrary case by adjusting the parameters in the more general joint unitary transformation without affecting the success probability. This fact strengthens and advances our knowledge that dissonance plays as a key ingredient in the assisted optimal state discrimination other than entanglement.

Let us discuss a necessary criterion for requiring dissonance in AOSD based on linear entropy. Initially, Alice prepares the system qubit in a mixed state $\rho = p_+|\psi_+\rangle\langle\psi_+| + p_-|\psi_-\rangle\langle\psi_-|$. She sends it to Bob and informs Bob that she would like to perform the AOSD, but she does not tell Bob anything about $p_\pm$ and $|\psi_\pm\rangle$. However, Bob may tell Alice that if the linear entropy of the state $\mathcal{S}_L(\rho) > 1/2$ (or the purity $\mathcal{P}(\rho) < 1/2$), then Alice needs dissonance for her assisted optimal state discrimination (see Fig. 3). Eventually, we would like to mention that local distinguishability of multipartite orthogonal quantum states was studied in [24], in which the local discrimination of entangled states also does not require any entanglement.

F.L.Z. is supported by NSF of China (Grant No. 11105097). J.L.C. is supported by National Basic Research Program (973 Program) of China under Grant No. 2012CB921900, NSF of China (Grant Nos. 10975075 and 11175089) and partly supported by National Research Foundation and Ministry of Education of Singapore.

---

[*] Electronic address: flzhang@tju.edu.cn

[†] Electronic address: chenjl@nankai.edu.cn

[1] R. Horodecki, P. Horodecki, M. Horodecki, K. Horodecki, Rev. Mod. Phys. **81**, 865(2009).
[2] J. S. Bell, Physics (Long Island City, N.Y.) **1**, 195 (1964).
[3] H. Ollivier and W. H. Zurek, Phys. Rev. Lett. **88**, 017901 (2001).
[4] L. Henderson and V. Vedral, J. Phys. A **34**, 6899 (2001).
[5] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, 2000).
[6] A. K. Ekert, Phys. Rev. Lett. **67**, 661 (1991).
[7] B. P. Lanyon, M. Barbieri, M. P. Almeida, and A. G. White, Phys. Rev. Lett. **101**, 200501 (2008).
[8] A. Datta, A. Shaji, and C. M. Caves, Phys. Rev. Lett. **100**, 050502 (2008).
[9] D. Cavalcanti, L. Aolita, S. Boixo, K. Modi, M. Piani, and A. Winter, Phys. Rev. A **83**, 032324 (2011).
[10] V. Madhok and A. Datta, Phys. Rev. A **83**, 032323 (2011).
[11] M. Piani, S. Gharibian, G. Adesso, J. Calsamiglia, P. Horodecki, and A. Winter, Phys. Rev. Lett. **106**, 220403 (2011).
[12] A. Streltsov, H. Kampermann, and D. Bruß, Phys. Rev. Lett. **106**, 160401 (2011).
[13] K. Modi, T. Paterek, W. Son, V. Vedral, and M. Williamson, Phys. Rev. Lett. **104**, 080501 (2010).
[14] L. Roa, J. C. Retamal, and M. Alid-Vaccarezza, Phys. Rev. Lett. **107**, 080401 (2011).
[15] J. V. Neumann, *Mathematical Foundations of Quantum Mechanics*, Vol. **2** (Princeton University Press, 1996).
[16] C. Helstrom, Journal of Statistical Physics **1**, 231 (1969).
[17] L. Roa, J. C. Retamal, and C. Saavedra, Phys. Rev. A **66**, 012103 (2002).
[18] M. A. Jafarizadeh, M. Rezaei, N. Karimi, and A. R. Amiri, Phys. Rev. A **77**, 042314 (2008).
[19] S. Pang and S. Wu, Phys. Rev. A **80**, 052320 (2009).
[20] When $\beta = \pi/4$ and $\delta = 0$, after performing a local unitary transformation $\mathcal{V} = \big(|+\rangle\langle 0| + |-\rangle\langle 1|\big) \otimes \mathbb{1}_a$ on Eq. (1), one can obtain the corresponding Eq. (1) in Ref. [14]. Here $|\pm\rangle = (|0\rangle \pm |1\rangle)/\sqrt{2}$.
[21] W. K. Wootters, Phys. Rev. Lett. **80**, 2245 (1998).
[22] M. Koashi and A. Winter, Phys. Rev. A **69**, 022309 (2004).
[23] V. Coffman, J. Kundu, and W. K. Wootters, Phys. Rev. A **61**, 052306 (2000).
[24] J. Walgate, A. J. Short, L. Hardy, and V. Vedral, Phys Rev Lett. **85**, 4972 (2000).